

Boundedness on inhomogeneous Lipschitz spaces of fractional integrals, singular integrals and hypersingular integrals associated to non-doubling measures.

A. Eduardo Gatto¹
DePaul University and CRM

To my sons

Abstract:

In the context of a finite measure metric space whose measure satisfies a growth condition, we prove "T1" type necessary and sufficient conditions for the boundedness of fractional integrals, singular integrals, and hypersingular integrals on inhomogeneous Lipschitz spaces. We also indicate how the results can be extended to the case of infinite measure. Finally we show applications to Real and Complex Analysis.

¹This paper was supported by a sabbatical leave from DePaul University and by a grant from the Ministerio de Educacion y Ciencia of Spain, Ref. # SAB2006-0118 for a quarter sabbatical visit to the Centre de Recerca Matemàtica in Barcelona.

The author wants to express a deep gratitude to these institutions, to the Analysis group of the Universidad Autònoma de Barcelona, in particular to Xavier Tolsa for the invitation and to the director and staff of the CRM for their pleasant hospitality and excellent resources.

Keywords: non-doubling measures, metric spaces, Lipschitz spaces, fractional integrals, singular integral, hypersingular integrals.

2000 MSC: 42B20, 26B35, 47B38, 47G10

1. Introduction. Definitions and Statement of the Theorems

Let (X, d, μ) be a finite measure metric space whose measure μ satisfies a n -dimensional growth condition, that is, (X, d) is a metric space and μ is a finite Borel measure that satisfies the following condition: there is $n > 0$ and a constant $A > 0$ such that $\mu(B_r) \leq Ar^n$, for all balls B_r of radius r and for all $r > 0$. Note that this condition allows the consideration of non-doubling as well as doubling measures.

Our results will apply to functions defined on the support of μ , of course the support of μ has to be well defined, where $\text{supp}(\mu)$ is the smallest closed set F such that for all Borel sets E , $E \subset F^c$, $\mu(E) = 0$. For example, if X is separable, then the support of μ is well defined. Furthermore to avoid any confusion we will assume that $X = \text{supp}(\mu)$.

The inhomogeneous Lipschitz-Hölder spaces of order β , $0 < \beta \leq 1$, will be denoted Λ_β and consists of all bounded functions f that satisfy

$\sup_{x \neq y \in X} \frac{|f(x) - f(y)|}{d^\beta(x, y)} < \infty$. The space Λ_β is a Banach space with the norm $\|f\|_{\Lambda_\beta} = \sup_{x \in X} |f(x)| + \sup_{x \neq y \in X} \frac{|f(x) - f(y)|}{d^\beta(x, y)}$. It will be useful to have a notation for each term in the norm, let $\sup(f) = \sup_{x \in X} |f(x)|$, and $|f|_\beta = \sup_{x \neq y \in X} \frac{|f(x) - f(y)|}{d^\beta(x, y)}$.

The results in this paper have extensions to the case $\mu(X) = \infty$, but the constants depend on the normalization of the integrals at infinity, we will indicate these extensions after the section on proofs. The letter C , c will denote constants not necessarily the same at each occurrence.

Let $\Omega = X \times X \setminus \Delta$, where $\Delta = \{(x, y) : x = y\}$. A function $L_\alpha(x, y) : \Omega \rightarrow \mathbb{C}$ will be called a standard fractional integral kernel of order α , $0 < \alpha < 1$, when there are constants B_1 and B_2 such that

$$(L1) \quad |L_\alpha(x, y)| \leq \frac{B_1}{d^{n-\alpha}(x, y)}.$$

$$(L2) \quad |L_\alpha(x_1, y) - L_\alpha(x_2, y)| \leq B_2 \frac{d^\gamma(x_1, x_2)}{d^{n-\alpha+\gamma}(x_1, y)}, \text{ for some } \gamma, \alpha < \gamma \leq 1, \text{ and } 2d(x_1, x_2) \leq d(x_1, y).$$

The fractional integral of order α of a function f in Λ_β is defined by:

$$L_\alpha f(x) = \int L_\alpha(x, y) f(y) d\mu(y).$$

Note that in particular $L_\alpha(x, y) = \frac{1}{d^{n-\alpha}(x, y)}$ is a standard fractional kernel of order α .

Theorem 1

Let $0 < \alpha < \gamma \leq 1$, $0 < \beta < 1$, and $\alpha + \beta \leq 1$ when $1 < n$ or $\alpha + \beta < n$ when $n \leq 1$. The following statements are equivalent:

- a) $L_\alpha 1 \in \Lambda_{\alpha+\beta}$.

b) $L_\alpha : \Lambda_\beta \rightarrow \Lambda_{\alpha+\beta}$ is bounded.

We define now the singular integral kernels that we will consider in Theorem 2 and Theorem 3. A function $K(x, y) : \Omega \rightarrow C$ will be called a standard singular integral kernel when there are constants C_1, C_2 and a number $\gamma, 0 < \gamma \leq 1$, such that

$$(S1) \quad |K(x, y)| \leq \frac{C_1}{d^n(x, y)}$$

$$(S2) \quad |K(x_1, y) - K(x_2, y)| \leq C_2 \frac{d^\gamma(x_1, x_2)}{d^{n+\gamma}(x_1, y)}, \text{ for } 2d(x_1, x_2) \leq d(x_1, y)$$

Let η be a function in $C^1[0, \infty)$ such that $0 \leq \eta \leq 1, \eta(s) = 0$ for $0 \leq s \leq 1/2$ and $\eta(s) = 1$ for $1 \leq s$. Let $K_\varepsilon(x, y) = \eta(\frac{d(x, y)}{\varepsilon})K(x, y)$, $\varepsilon > 0$ where $K(x, y)$ is a standard singular integral kernel. We will denote T_ε the operator $T_\varepsilon f(x) = \int K_\varepsilon(x, y)f(y)d\mu(y)$.

Theorem 2

Let $K(x, y)$ be a standard singular integral kernel. Let $0 < \beta < \min(n, \gamma)$. The following two statements are equivalent:

a) $\|T_\varepsilon 1\|_{\Lambda_\beta} \leq C$, for all $\varepsilon > 0$.

b) $T_\varepsilon : \Lambda_\beta \rightarrow \Lambda_\beta$ are bounded and $\|T_\varepsilon\|_{\Lambda_\beta \rightarrow \Lambda_\beta} \leq C'$, for all $\varepsilon > 0$.

One of the novelties in this Theorem is that the cancellation condition (S3) for all x (see below) follows from part b).

In Theorem 3 we will consider Principal Value Singular Integrals. We will denote by Lip_β the space of classes of measurable functions f for which there is a $g \in \Lambda_\beta$ such that $f = g$ except for a set E that depends on f , with $\mu(E) = 0$. The norm of f in Lip_β is defined as $\|f\|_{Lip_\beta} = \|f\|_\infty + |f|_\beta$, where $|f|_\beta = \sup_{x \neq y \in X} \frac{|g(x) - g(y)|}{d^\beta(x, y)} = \sup_{x \neq y \in X - E} \frac{|f(x) - f(y)|}{d^\beta(x, y)}$.

We also need to add the following two conditions on the kernel:

$$(S3) \quad \left| \int_{r_1 < d(x, y) < r_2} K(x, y)d\mu(y) \right| \leq C_3 \text{ for all } 0 < r_1 < r_2 < \infty, \mu - a.e \text{ in } x.$$

$$(S4) \quad \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x, y) < 1} K(x, y)d\mu(y) \text{ exists } \mu - a.e \text{ in } x.$$

The principal value singular integral of a function $f \in Lip_\beta$ is defined by

$$Kf(x) = \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x, y)} K(x, y)f(y)d\mu(y)$$

Theorem 3

Let $K(x, y)$ be a standard singular integral kernel that in addition satisfies (S3) and (S4). Let $0 < \beta < \min(n, \gamma)$ and $f \in Lip_\beta$. Then $Kf(x)$ is well defined $\mu - a.e.$ and the following two statements are equivalent:

a) $K1 \in Lip_\beta$

b) $K : Lip_\beta \rightarrow Lip_\beta$ is bounded

A function $D_\alpha(x, y) : \Omega \rightarrow C$ will be called a standard hypersingular kernel of order α , $0 < \alpha < 1$, when there are constants E_1 and E_2 such that:

$$(D1) \quad |D_\alpha(x, y)| \leq \frac{E_1}{d^{n+\alpha}(x, y)},$$

$$(D2) \quad |D_\alpha(x_1, y) - D_\alpha(x_2, y)| \leq E_2 \frac{d^\gamma(x_1, x_2)}{d^{n+\alpha+\gamma}(x_1, y)}, \text{ for some } \gamma, 0 < \gamma \leq 1, \text{ and } 2d(x_1, x_2) \leq d(x_1, y).$$

The hypersingular integral of order α of a function $f \in \Lambda_\beta$ $\alpha < \beta \leq 1$ is defined by:

$$D^\alpha f(x) = \int D_\alpha(x, y) [f(y) - f(x)] d\mu(y)$$

Note that in particular $D_\alpha(x, y) = \frac{1}{d^{n+\alpha}(x, y)}$ is a standard hypersingular kernel of order α when $X = R^n$ and μ is the Lebesgue measure, and we have $\int \frac{1}{d^{n+\alpha}(x, y)} [f(y) - f(x)] dy = c_\alpha (\Delta^{\frac{\alpha}{2}} f)(x)$ for f sufficiently smooth and $0 < \alpha < 2$. [S]

Theorem 4

Let $0 < \alpha < \beta \leq 1$ and $\beta - \alpha < n$. Then $D^\alpha : \Lambda_\beta \rightarrow \Lambda_{\beta-\alpha}$ is bounded.

Note that $D^\alpha 1 = 0$ by definition. Also, Theorem 4 and its proof are valid without changes in the case $\mu(X) = \infty$.

2. Proofs

We would like to point out that the proofs are based on classical methods, see for example [Z], adjusted to the modern "T1" formulation and to the present general context. For carrying out the proofs we need the following known lemma about measures that satisfy the n -dimensional growth condition.

Lemma

Let (X, d, μ) be a measure metric space such that μ satisfies the n -dimensional growth condition, $r > 0$. Then

1. $\int_{d(x, y) < r} \frac{1}{d^{n-\delta}(x, y)} d\mu(y) \leq c_1 r^\delta, \quad 0 < \delta < n.$
2. $\int_{r \leq d(x, y)} \frac{1}{d^{n+\delta}(x, y)} d\mu(y) \leq c_2 r^{-\delta}, \quad 0 < \delta$
3. $\int_{r/2 \leq d(x, y) < r} \frac{1}{d^n(x, y)} d\mu(y) \leq c_3$

Proof of the Lemma

The three parts are a consequence of the growth condition. To prove part1, we rewrite the integral as a series and mayorize each term using the growth condition and we add the resulting series. In detail we have:

$$\begin{aligned} \int_{d(x, x_o) < r} \frac{1}{d^{n-\delta}(x, x_o)} d\mu(x) &= \sum_{k=0}^{\infty} \int_{2^{-k-1}r \leq d(x, x_o) < 2^{-k}r} \frac{1}{d^{n-\delta}(x, x_o)} d\mu(x) \leq \\ &\sum_{k=0}^{\infty} \frac{\mu(B_{2^{-k-1}r}(x_o))}{(2^{-k-1}r)^{n-\delta}} \leq A \sum_{k=0}^{\infty} \frac{(2^{-k}r)^n}{(2^{-k-1}r)^{n-\delta}} = Ar^{\delta} \left(\frac{2^n}{2^{\delta}-1} \right). \end{aligned}$$

To prove part 2 we perform a similar estimate:

$$\begin{aligned} \int_{d(x, x_o) \geq r} \frac{1}{d^{n+\delta}(x, x_o)} d\mu(x) &= \sum_{k=0}^{\infty} \int_{2^k r \leq d(x, x_o) < 2^{k+1}r} \frac{1}{d^{n+\delta}(x, x_o)} d\mu(x) \leq \\ &\sum_{k=0}^{\infty} \frac{\mu(B_{2^{k+1}r}(x_o))}{(2^{k+1}r)^{n+\delta}} \leq A \sum_{k=0}^{\infty} \frac{(2^{k+1}r)^n}{(2^{k+1}r)^{n+\delta}} = Ar^{-\delta} \left(\frac{2^n 2^{\delta}}{2^{\delta}-1} \right) \end{aligned}$$

Finally for part 3 we have:

$$\int_{r/2 \leq d(x, y) < r} \frac{1}{d^n(x, y)} d\mu(y) \leq \frac{\mu(B_r(x_o))}{(r/2)^n} \leq A2^n.$$

Proof of Theorem 1

Observe first that $1 \in \Lambda_{\beta}$ and therefore condition b) implies condition a). We will prove now that condition a) implies condition b). We can just consider the case $L_{\alpha}(x, y) = \frac{1}{d^{n-\alpha}(x, y)}$, because the general case is proven in the same way, and we will denote $L_{\alpha} = I_{\alpha}$.

Condition (L1) is clearly valid. To show that condition (L2) is verified, we use the Mean Value Theorem. Consider $2d(x_1, x_2) \leq d(x_1, y)$, and $0 < \theta < 1$ we have:

$$\begin{aligned} \left| \frac{1}{d^{n-\alpha}(x_1, y)} - \frac{1}{d^{n-\alpha}(x_2, y)} \right| &\leq \sup_{\theta} |(-n + \alpha)(\theta d(x_1, y) + (1 - \theta)d(x_2, y))^{-n+\alpha-1}| \cdot \\ &|d(x_1, y) - d(x_2, y)| \leq B_2 \frac{d(x_1, x_2)}{d^{n-\alpha+1}(x_1, y)} \end{aligned}$$

Now we will estimate $\sup(I_{\alpha}f)$. Let $x \in X$. We will use the Lemma to obtain

$$|I_{\alpha}f(x)| \leq \int_{d(x, y) < 1} \frac{|f(y)|}{d^{n-\alpha}(x, y)} d\mu(y) + \int_{1 \leq d(x, y)} \frac{|f(y)|}{d^{n-\alpha}(x, y)} d\mu(y) \leq \|f\|_{\infty} (c_1 + \mu(X)),$$

and therefore $\sup(I_\alpha f) \leq \sup(f)(c_1 + \mu(X))$. We will estimate next $|I_\alpha f|_{(\beta)}$. We write

$$\begin{aligned} I_\alpha f(x_1) - I_\alpha f(x_2) &= \int_X \frac{f(y)}{d^{n-\alpha}(x_1, y)} d\mu(y) - \int_X \frac{f(y)}{d^{n-\alpha}(x_2, y)} d\mu(y) = \int_X \frac{f(y) - f(x_1)}{d^{n-\alpha}(x_1, y)} d\mu(y) + \\ &f(x_1) \int_X \frac{1}{d^{n-\alpha}(x_1, y)} d\mu(y) - \int_X \frac{f(y) - f(x_1)}{d^{n-\alpha}(x_2, y)} d\mu(y) - f(x_1) \int_X \frac{1}{d^{n-\alpha}(x_2, y)} d\mu(y) = \\ &\int_X \frac{f(y) - f(x_1)}{d^{n-\alpha}(x_1, y)} d\mu(y) - \int_X \frac{f(y) - f(x_1)}{d^{n-\alpha}(x_2, y)} d\mu(y) + f(x_1) [I_\alpha 1(x_1) - I_\alpha 1(x_2)]. \end{aligned}$$

The last term can be majorized using the hypothesis, and we have $|f(x_1) [I_\alpha 1(x_1) - I_\alpha 1(x_2)]| \leq c \sup(f) d^{\alpha+\beta}(x_1, x_2)$.

Let now $r = d(x_1, x_2)$ and $B_{2r}(x_1)$ the ball of radius $2r$ and center x_1 . We write

$$\begin{aligned} &\left| \int_X \frac{f(y) - f(x_1)}{d^{n-\alpha}(x_1, y)} d\mu(y) - \int_X \frac{f(y) - f(x_1)}{d^{n-\alpha}(x_2, y)} d\mu(y) \right| \leq \\ &\int_{B_{2r}(x_1)} \frac{|f(y) - f(x_1)|}{d^{n-\alpha}(x_1, y)} d\mu(y) + \int_{B_{2r}(x_1)} \frac{|f(y) - f(x_1)|}{d^{n-\alpha}(x_2, y)} d\mu(y) + \\ &\int_{B_{2r}^c(x_1)} |f(y) - f(x_1)| \left| \frac{1}{d^{n-\alpha}(x_1, y)} - \frac{1}{d^{n-\alpha}(x_2, y)} \right| d\mu(y) = J_1 + J_2 + J_3 \end{aligned}$$

For the first term using the lemma we have

$$J_1 \leq |f|_{(\beta)} \int_{B_{2r}(x_1)} \frac{d^\beta(x_1, y)}{d^{n-\alpha}(x_1, y)} d\mu(y) \leq c |f|_{(\beta)} r^{\alpha+\beta} = c |f|_{(\beta)} d^{\alpha+\beta}(x_1, x_2).$$

For the second term we write

$$J_2 \leq |f|_{(\beta)} \int_{B_{3r}(x_2)} \frac{2r}{d^{n-\alpha}(x_2, y)} d\mu(y) \leq c |f|_{(\beta)} d^{\alpha+\beta}(x_1, x_2),$$

For the third term we use (L2) and the lemma to get

$$J_3 \leq |f|_{(\beta)} \int_{B_{2r}^c(x_1)} \frac{B_2}{d^{n-\alpha-\beta}(x_1, y)} d\mu(y) \leq c |f|_{(\beta)} d^{\alpha+\beta}(x_1, x_2)$$

Collecting the previous estimates, we have

$$\|I_\alpha f\|_{\Lambda_\beta} \leq C \|f\|_{\Lambda_\beta}$$

This concludes the proof of Theorem 1.

Proof of Theorem 2

Observe first that $1 \in \Lambda_\beta$ and therefore condition b) implies condition a).

Before doing the proof of the theorem and for the sake of completeness, we will show that K_ε satisfies conditions (S1) and (S2) with constants independent of ε .

Condition (S1) is true because η is bounded. To show condition (S2), assume that $2d(x_1, x_2) \leq d(x_1, y)$ and consider the following two cases:

Case 1: $1 < \frac{d(x_1, y)}{\varepsilon}$ and $1 < \frac{d(x_2, y)}{\varepsilon}$. In this case $K_\varepsilon(x, y) = K(x, y)$, and therefore (S2) is true with the same constant.

Case 2: $1 \geq \frac{d(x_1, y)}{\varepsilon}$ or $1 \geq \frac{d(x_2, y)}{\varepsilon}$. Assume $1 > \frac{d(x_1, y)}{\varepsilon}$.

We write

$$|K_\varepsilon(x_1, y) - K_\varepsilon(x_2, y)| \leq \left| \eta\left(\frac{d(x_1, y)}{\varepsilon}\right) - \eta\left(\frac{d(x_2, y)}{\varepsilon}\right) \right| |K(x_1, y)| + \left| \eta\left(\frac{d(x_2, y)}{\varepsilon}\right) \right| |K(x_1, y) - K(x_2, y)|$$

The first term above is less than or equal to

$$\begin{aligned} \|\eta'\|_\infty \frac{|d(x_1, y) - d(x_2, y)|}{\varepsilon} |K(x_1, y)| &\leq \|\eta'\|_\infty \frac{d(x_1, x_2)}{\varepsilon} |K(x_1, y)| \leq \\ &c \left(\frac{d(x_1, x_2)}{\varepsilon}\right)^\gamma |K(x_1, y)| \leq c \frac{d^\gamma(x_1, x_2)}{d^{n+\gamma}(x_1, y)} \end{aligned}$$

On the other hand the second term is less than or equal to $c |K(x_1, y) - K(x_2, y)| \leq c \frac{d^\gamma(x_1, x_2)}{d^{n+\gamma}(x_1, y)}$. If $1 \geq \frac{d(x_2, y)}{\varepsilon}$ the proof is similar.

To show that condition a) implies condition b), the first step is to get the cancellation (S3) of the kernel, for all $x \in X$.

Observe that for $0 < r_1 < r_2 < \infty$, we have

$$\begin{aligned} T_{r_1} 1(x) - T_{r_2} 1(x) &= \int_{\frac{1}{2}r_1 < d(x, y) \leq r_1} \eta\left(\frac{d(x, y)}{r_1}\right) K(x, y) d\mu(y) + \int_{r_1 < d(x, y)} K(x, y) d\mu(y) \\ &\quad - \int_{\frac{1}{2}r_2 < d(x, y) < r_2} \eta\left(\frac{d(x, y)}{r_2}\right) K(x, y) d\mu(y) - \int_{r_2 \leq d(x, y)} K(x, y) d\mu(y) \end{aligned}$$

Since the left hand side is uniformly bounded in r and x , and also the first and third terms are uniformly bounded because of the growth condition (see lemma), it follows that

$$(S3) \quad \left| \int_{r_1 < d(x,y) < r_2} K(x,y) d\mu(y) \right| \leq C, \text{ for all } x.$$

Now, we will estimate $\sup |T_\varepsilon f(x)|$. Observe first that

$$\begin{aligned} T_\varepsilon f(x) &= \int_{d(x,y) \leq 1} K_\varepsilon(x,y) f(y) d\mu(y) + \int_{d(x,y) > 1} K_\varepsilon(x,y) f(y) d\mu(y) = \\ &= \int_{d(x,y) \leq 1} K_\varepsilon(x,y) f(y) - f(x) d\mu(y) + f(x) \int_{\frac{1}{2}\varepsilon < d(x,y) \leq \varepsilon} K_\varepsilon(x,y) d\mu(y) \\ &\quad + f(x) \int_{\varepsilon < d(x,y) \leq 1} K_\varepsilon(x,y) d\mu(y) + \int_{d(x,y) > 1} K_\varepsilon(x,y) f(y) d\mu(y) \end{aligned}$$

Now, by conditions (S1), (S3) and the growth condition we can bound the absolute value of the terms above by $\|f\|_{\Lambda_\beta}$ and therefore $\sup_{x \in X} |T_\varepsilon f(x)| \leq c \|f\|_{\Lambda_\beta}$.

Next, we estimate $\sup_{x \neq y} \frac{|T_\varepsilon f(x) - T_\varepsilon f(y)|}{d^\beta(x,y)}$. We consider the difference $T_\varepsilon f(x_1) - T_\varepsilon f(x_2)$, and the following decomposition:

$$\begin{aligned} T_\varepsilon f(x_1) - T_\varepsilon f(x_2) &= \int K_\varepsilon(x_1, y) f(y) d\mu(y) - \int K_\varepsilon(x_2, y) f(y) d\mu(y) \\ &= \int K_\varepsilon(x_1, y) [f(y) - f(x_1)] d\mu(y) + f(x_1) \int K_\varepsilon(x_1, y) d\mu(y) - \\ &\quad \int K_\varepsilon(x_2, y) [f(y) - f(x_1)] d\mu(y) - f(x_1) \int K_\varepsilon(x_2, y) d\mu(y) = \\ &= \int K_\varepsilon(x_1, y) [f(y) - f(x_1)] d\mu(y) + \int K_\varepsilon(x_2, y) [f(y) - f(x_1)] d\mu(y) + \\ &\quad f(x_1) [T_\varepsilon 1(x_1) - T_\varepsilon 1(x_2)] \end{aligned}$$

Observe now that the last term can be estimated using the hypothesis and we have

$$|f(x_1) [T_\varepsilon 1(x_1) - T_\varepsilon 1(x_2)]| \leq c \sup(f) d^\beta(x_1, x_2).$$

To estimate the first two terms, let $r = d(x_1, x_2)$, we rewrite them as follows:

$$\begin{aligned} & \int K_\varepsilon(x_1, y) [f(y) - f(x_1)] d\mu(y) + \int K_\varepsilon(x_2, y) [f(y) - f(x_1)] d\mu(y) = \\ & \int_{d(x_1, y) < 3r} K_\varepsilon(x_1, y) [f(y) - f(x_1)] d\mu(y) + \int_{d(x_1, y) < 3r} K_\varepsilon(x_2, y) [f(y) - f(x_1)] d\mu(y) + \\ & \int_{3r < d(x_1, y)} [f(y) - f(x_1)] [K_\varepsilon(x_1, y) - K_\varepsilon(x_2, y)] d\mu(y) = H_1 + H_2 + H_3 \end{aligned}$$

The absolute value of H_3 can be estimated as follows,

$$|H_3| \leq |f|_\beta d^\gamma(x_1, x_2) \int_{3r < d(x_1, y)} \frac{d^\beta(x_1, y)}{d^{n+\gamma}(x_1, y)} d\mu(y) \leq c |f|_\beta d^\beta(x_1, x_2)$$

For $|H_1|$ we have

$$|H_1| \leq |f|_{[\beta]} \int_{d(x_1, y) < 3r} \frac{C_1}{d^{n-\beta}(x_1, y)} d\mu(y) \leq c |f|_{[\beta]} d^\beta(x_1, x_2)$$

Finally to estimate H_2 we write

$$\begin{aligned} & \int_{d(x_1, y) < 3r} K_\varepsilon(x_2, y) [f(y) - f(x_1)] d\mu(y) = \\ & \int_{d(x_1, y) < 3r} K_\varepsilon(x_2, y) [f(y) - f(x_2)] d\mu(y) \\ & + [f(x_2) - f(x_1)] \int_{\{y: \varepsilon/2 < d(x_2, y)\} \cap \{y: d(x_1, y) < 3r\}} K_\varepsilon(x_2, y) d\mu(y) = J_1 + J_2 \end{aligned}$$

For the first term we have

$$|J_1| \leq \int_{d(x_2, y) < 4r} \frac{c \|f\|_{(\beta)}}{d^{n-\beta}(x_2, y)} d\mu(y) \leq c |f|_\beta d^\beta(x_1, x_2)$$

To estimate J_2 consider first

$$\int_{d(x_1, y) < 3r} K_\varepsilon(x_2, y) d\mu(y) = \int_{d(x_2, y) < 2r} K_\varepsilon(x_2, y) d\mu(y) +$$

$$\int_{\{y:d(x_1,y)<3r\}\setminus\{y:d(x_2,y)<2r\}} K_\varepsilon(x_2,y)d\mu(y)$$

Observe now that

$$\left| \int_{d(x_2,y)<2r} K_\varepsilon(x_2,y)d\mu(y) \right| \leq C_3$$

and using part 3 of the lemma we get

$$\left| \int_{\{y:d(x_1,y)<3r\}\setminus\{y:d(x_2,y)<2r\}} K_\varepsilon(x_2,y)d\mu(y) \right| \leq \int_{\{y:2r<d(x_2,y)<4r\}} |K_\varepsilon(x_2,y)| d\mu(y) \leq c$$

therefore

$$|J_2| \leq c \|f\|_\beta d^\beta(x_1, x_2)$$

collecting the estimates we have:

$$|K_\varepsilon f(x_1) - K_\varepsilon f(x_2)| \leq c \|f\|_{\Lambda_\beta} d^\beta(x_1, x_2)$$

and finally

$$\|K_\varepsilon f\|_{\Lambda_\beta} \leq c \|f\|_{\Lambda_\beta},$$

with c independent of ε .

Proof of Theorem 3

Observe first that $1 \in Lip_\beta$ and therefore condition b) implies condition a). Let $f \in Lip_\beta$, we will show that

$$Kf(x) = \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x,y)} K(x,y)f(y)d\mu(y)$$

exists $\mu - a.e.$ Assume $\varepsilon < 1$, we can write

$$\begin{aligned} Kf(x) &= \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x,y) < 1} K(x,y) [f(y) - f(x)] d\mu(y) + \\ &f(x) \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x,y) < 1} K(x,y) d\mu(y) + \int_{1 \leq d(x,y)} K(x,y) f(y) d\mu(y). \end{aligned}$$

Note that the first integral converges absolutely, the limit of the second term exists by hypothesis and finally last integral converges absolutely because the integrand is bounded. Furthermore, we have $\|Kf\|_\infty \leq c \|f\|_{Lip_\beta}$.

We will estimate now $Kf(x_1) - Kf(x_2)$ for x_1, x_2 two points for which $Kf(x)$ exists. This part of the proof is very similar to the same part in Theorem 2. We write

$$\begin{aligned}
Kf(x_1) - Kf(x_2) &= \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_1, y)} K(x_1, y) f(y) d\mu(y) - \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y)} K(x_2, y) f(y) d\mu(y) \\
&= \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_1, y)} K(x_1, y) [f(y) - f(x_1)] d\mu(y) + f(x_1) \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_1, y)} K(x_1, y) d\mu(y) - \\
&\quad \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y)} K(x_2, y) [f(y) - f(x_1)] d\mu(y) - f(x_1) \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y)} K(x_2, y) d\mu(y) = \\
&\quad \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_1, y)} K(x_1, y) [f(y) - f(x_1)] d\mu(y) + \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y)} K(x_2, y) [f(y) - f(x_1)] d\mu(y) + \\
&\quad f(x_1) [K1(x_1) - K1(x_2)]
\end{aligned}$$

Observe now that the last term can be estimated using the hypothesis and we have

$$|f(x_1) [K1(x_1) - K1(x_2)]| \leq c \|f\|_{\infty} d^{\beta}(x_1, x_2).$$

To estimate the first two terms, let $r = d(x_1, x_2)$, and $\varepsilon < r$, we rewrite them as follows:

$$\begin{aligned}
&\lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_1, y)} K(x_1, y) [f(y) - f(x_1)] d\mu(y) + \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y)} K(x_2, y) [f(y) - f(x_1)] d\mu(y) = \\
&\quad \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_1, y) < 3r} K(x_1, y) [f(y) - f(x_1)] d\mu(y) + \\
&\quad \lim_{\varepsilon \rightarrow 0} \int_{\{y: \varepsilon < d(x_2, y)\} \cap \{y: d(x_1, y) < 3r\}} K(x_2, y) [f(y) - f(x_1)] d\mu(y) + \\
&\quad \lim_{\varepsilon \rightarrow 0} \int_{3r < d(x_1, y)} [f(y) - f(x_1)] [K(x_1, y) - K(x_2, y)] d\mu(y) = H_1 + H_2 + H_3
\end{aligned}$$

The absolute value of H_3 can be estimated as follows,

$$|H_3| \leq |f|_{\beta} d^{\gamma}(x_1, x_2) \int_{3r < d(x_1, y)} \frac{d^{\beta}(x_1, y)}{d^{n+\gamma}(x_1, y)} d\mu(y) \leq c |f|_{\beta} d^{\beta}(x_1, x_2)$$

For $|H_1|$ we have

$$|H_1| \leq |f|_\beta \int_{d(x_1, y) < 3r} \frac{C_1}{d^{n-\beta}(x_1, y)} d\mu(y) \leq c |f|_\beta d^\beta(x_1, x_2)$$

Finally to estimate H_2 we write

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \int_{\{y: \varepsilon < d(x_2, y)\} \cap \{y: d(x_1, y) < 3r\}} K(x_2, y) [f(y) - f(x_1)] d\mu(y) = \\ & \lim_{\varepsilon \rightarrow 0} \int_{\{y: \varepsilon < d(x_2, y)\} \cap \{y: d(x_1, y) < 3r\}} K(x_2, y) [f(y) - f(x_2)] d\mu(y) \\ & + [f(x_2) - f(x_1)] \lim_{\varepsilon \rightarrow 0} \int_{\{y: \varepsilon < d(x_2, y)\} \cap \{y: d(x_1, y) < 3r\}} K(x_2, y) d\mu(y) = J_1 + J_2 \end{aligned}$$

For the first term we have

$$|J_1| \leq \int_{d(x_2, y) < 4r} \frac{c \|f\|_{(\beta)}}{d^{n-\beta}(x_2, y)} d\mu(y) \leq c |f|_\beta d^\beta(x_1, x_2)$$

To estimate the second J_2 consider first

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{\{y: \varepsilon < d(x_2, y)\} \cap \{y: d(x_1, y) < 3r\}} K(x_2, y) d\mu(y) &= \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y) < 2r} K(x_2, y) d\mu(y) + \\ & \int_{\{y: d(x_1, y) < 3r\} \setminus \{y: d(x_2, y) < 2r\}} K(x_2, y) d\mu(y) \end{aligned}$$

Observe now that

$$\left| \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x_2, y) < 2r} K(x_2, y) d\mu(y) \right| \leq C_3$$

and using part 3 of the lemma we get

$$\left| \int_{\{y: d(x_1, y) < 3r\} \setminus \{y: d(x_2, y) < 2r\}} K(x_2, y) d\mu(y) \right| \leq \int_{\{y: 2r < d(x_2, y) < 4r\}} |K(x_2, y)| d\mu(y) \leq c$$

therefore

$$|J_2| \leq c |f|_\beta d^\beta(x_1, x_2)$$

collecting the estimates we have:

$$|Kf(x_1) - Kf(x_2)| \leq c \|f\|_{Lip_\beta} d^\beta(x_1, x_2)$$

and finally

$$\|Kf\|_{Lip_\beta} \leq c \|f\|_{Lip_\beta}$$

This concludes the proof of Theorem 3.

Proof of Theorem 4

We will prove the theorem for $D_\alpha(x, y) = \frac{1}{d^{n+\alpha}(x, y)}$, the general case is identical. Note that the proof is also valid for $\mu(X) = \infty$.

We will estimate first $\sup(D_\alpha f)$ for $f \in \Lambda_\beta$. We use part 2 of the Lemma to write

$$|D^\alpha f(x)| \leq \int_{d(x, y) \leq 1} \frac{|f(y) - f(x)|}{d^{n+\alpha}(x, y)} d\mu(y) + c \sup(f) \leq |f|_\beta \int_{d(x, y) \leq 1} \frac{1}{d^{n+\alpha-\beta}(x, y)} d\mu(y) + c \sup(f)$$

Since $0 < \alpha < \beta \leq 1$, we use part 1 of the lemma to estimate the integral and we obtain that $D^\alpha f(x)$ is well defined everywhere and

$$\sup(D^\alpha f) \leq c \|f\|_{\Lambda_\beta}.$$

To estimate $|D^\alpha f|_\alpha$, we consider $r = d(x, y)$ and write

$$\begin{aligned} D^\alpha f(x_1) - D^\alpha f(x_2) &= \int_{d(x_1, y) \leq 2r} \frac{f(y) - f(x_1)}{d^{n+\alpha}(x_1, y)} d\mu(y) - \int_{d(x_1, y) \leq 2r} \frac{f(y) - f(x_2)}{d^{n+\alpha}(x_2, y)} d\mu(y) + \\ &+ \int_{d(x_1, y) > 2r} [f(y) - f(x_1)] \left[\frac{1}{d^{n+\alpha}(x_1, y)} - \frac{1}{d^{n+\alpha}(x_2, y)} \right] d\mu(y) - \\ &\int_{d(x_1, y) > 2r} \frac{f(x_1) - f(x_2)}{d^{n+\alpha}(x_2, y)} d\mu(y) \end{aligned}$$

Using part 1 of the Lemma and the fact that f is in Λ_β we can obtain that each of the first two terms converges absolutely and is bounded by $c |f|_\beta d^{\beta-\alpha}(x_1, x_2)$. Using part 2 of the Lemma we can also obtain that the fourth term converges absolutely and is bounded by $c |f|_\beta d^{\beta-\alpha}(x_1, x_2)$.

To estimate the third term observe first that for $2d(x_1, x_2) \leq d(x_1, y)$,

$$\left| \frac{1}{d^{n+\alpha}(x_1, y)} - \frac{1}{d^{n+\alpha}(x_2, y)} \right| \leq \sup_\theta |(-n - \alpha)(\theta d(x_1, y) + (1 - \theta)d(x_2, y))^{-n-\alpha-1}|.$$

$$|d(x_1, y) - d(x_2, y)| \leq c \frac{d(x_1, x_2)}{d^{n+\alpha+1}(x_1, y)}.$$

Therefore using this estimate, the fact that $f \in \Lambda_\beta$ and the part 2 of Lemma we obtain that the third term converges absolutely and is less than or equal to $c |f|_\beta d^{\beta-\alpha}(x_1, x_2)$ and consequently $|D^\alpha|_{(\beta-\alpha)} \leq c |f|_\beta$.

Finally combining the two estimates we get $\|D^\alpha f\|_{\Lambda_{\beta-\alpha}} \leq c \|f\|_{\Lambda_\beta}$.

To extend Theorem 1 and Theorem 3, the fractional integrals and singular integrals have to be redefined so they converge for $d(x, y) > 1$. The operator's norm in each result will depend on the normalization. We will denote with ' ' the normalizations. Let $x_o \in X$ be a fixed point for which (S4) is valid and define:

$$L'_\alpha f(x) = \int [L_\alpha(x, y) - L_\alpha(x_o, y)] f(y) d\mu(y)$$

$$K' f(x) = \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < d(x, y)} [K(x, y) - K(x_o, y)] f(y) d\mu(y)$$

Applications

In this section we will illustrate some applications of the theorems. I am indebted to Joaquim Bruna for pointing out to me the Theorem of Mark Krein and to Joan Verdera for several generous discussions on applications 1 and 2.

1. The purpose of this application is to obtain boundedness in L^2 of some singular integrals in the context of non-doubling measure metric spaces of finite measure. Following [T], a singular integral associated to μ is said to be bounded in L^2 when there is a constant C such that $\|K_\varepsilon f\|_{L^2} \leq C \|f\|_{L^2}$, for all $\varepsilon > 0$, where $K_\varepsilon f(x) = \int_{d(x, y) > \varepsilon} K(x, y) f(y) d\mu(y)$. We will apply Theorem 2 and the following Theorem of Mark Krein (see[FMM] for its proof and application to the classical case, and [W] for the case of spaces of homogeneous type).

M. Krein's Theorem:

Let H be a real or complex Hilbert space with inner product $(\cdot, \cdot)$ and norm $\|\cdot\|_H$. Let $D \subset H$ be a Banach space dense in H and such that $\|x\|_H \leq C \|x\|_D$ for $x \in D$. Let A and B be two linear operator such that $\|Ax\|_D \leq C_A \|x\|_D$, $\|Bx\|_D \leq C_B \|x\|_D$, $x \in D$ and $(Ax, y) = (x, By)$ for all $x, y \in D$. Then $\|Ax\|_H \leq (C_A C_B)^{\frac{1}{2}} \|x\|_H$, $\|Bx\|_H \leq (C_A C_B)^{\frac{1}{2}} \|x\|_H$, $x \in D$, and both extend to bounded operator on H .

In our application, we will consider $H = L^2$ and $D = \Lambda_\beta$. Since X has finite measure we clearly have $\|f\|_{L^2} \leq \mu(X)^{\frac{1}{2}} \|f\|_{\Lambda_\beta}$, but we need the extra assumption Λ_β dense in L^2 . Let now $K(x, y)$ be a standard singular integral kernel and $K^*(x, y) = \overline{K(y, x)}$. Assume that $K^*(x, y)$ also satisfies (S2). Let $A = T_\varepsilon$ and $B = T_\varepsilon^*$ the corresponding smooth truncations. If $\|T_\varepsilon 1\|_{\Lambda_\beta} \leq C'$ and $\|T_\varepsilon^* 1\|_{\Lambda_\beta} \leq C''$ for all $\varepsilon > 0$, then by Theorem 2 and Krein's Theorem there is C such that $\|T_\varepsilon f\|_{L^2} \leq C \|f\|_{L^2}$, $f \in \Lambda_\beta$, for all $\varepsilon > 0$. Consequently $\|K_\varepsilon f\|_{L^2} \leq C \|f\|_{L^2}$, $f \in \Lambda_\beta$ for all $\varepsilon > 0$, and it extends to a bounded operator in L^2 , same conclusion for K^* . In addition, Nazarov, Treil, and Volberg have extended the classical result of Calderon-Zygmund on the boundedness in L^p , $1 < p < \infty$, of

singular integrals bounded in L^2 , to non-doubling separable measure metric spaces, see [NTV].

2. The second application has appeared in [MOV]. In this paper the authors need to study the boundedness properties of the Restricted Beurling Transform, $B_\Omega f = B(f\chi_\Omega)$, on $Lip_\varepsilon(\Omega)$ where Ω is a bounded domain in R^n with boundary of class $C^{1+\varepsilon}$, $0 < \varepsilon < 1$. Mateu, Orobitg and Verdera prove the following more general result: "Let Ω be a bounded domain with boundary of class $C^{1+\varepsilon}$, $0 < \varepsilon < 1$, and let T be an even smooth homogeneous Calderon-Zygmund operator. Then T_Ω maps $Lip_\varepsilon(\Omega)$ into $Lip_\varepsilon(\Omega)$ and also $Lip_\varepsilon(\Omega)$ into $Lip_\varepsilon(\Omega^c)$ ". Their proof, which is non-trivial, consists in showing that condition (S3) and part a) of Theorem 3 above are met.

3. The third application is related to M. Riesz Fractional Calculus associated to non-doubling measures. Applying Theorem 1 and Theorem 4 we can obtain that the composition of a Riesz fractional integral $I_\alpha f(x) = \int \frac{1}{d^{n-\alpha}(x,y)} f(y) d\mu(y)$ and a fractional derivative $D^\alpha f(x) = \int \frac{[f(y)-f(x)]}{d^{n+\alpha}(x,y)} d\mu(y)$ of the same order $D^\alpha I_\alpha$, as well as its transpose $I_\alpha D^\alpha$, are bounded on Λ_β , when $I_\alpha 1 \in \Lambda_{\alpha+\beta}$, $\alpha + \beta < 1$. In addition, it was shown in [G] that these composition are singular integral operators associated to μ

References

- [FMM] E. Fabes, I. Mitrea and M. Mitrea, "On the Boundedness of Singular Integrals". Pacific Journal of Mathematics, Vol. 189, #1, 1999.
- [GG1] J. Garcia-Cuerva and A. Eduardo Gatto, "Boundedness properties of fractional integrals operators associated to non-doubling measures". Studia Mathematica 162 (3) (2004).
- [GG2] J. Garcia-Cuerva and A. Eduardo Gatto, "Lipschitz Spaces and Calderon-Zygmund operators associated to non-doubling measures". Publ.Mat. 49 (2005), 285-296.
- [G] A. Eduardo Gatto, "On fractional calculus associated to doubling and non-doubling measures". Harmonic Analysis Calderon-Zygmund and Beyond .Contemporary Mathematics, # 411, Amer. Mat. Soc. 2006
- [GV] A. Eduardo Gatto and Steven Vàgi, "On molecules and Fractional Integrals on Spaces of Homogeneous Type with finite measure". Studia Mathematica, Vol. 103 (1), (1992), pp. 25-39
- [MOV] J. Mateu, J. Orobitg, and J. Verdera. "Extra cancellation of even Calderon-Zygmund operators and Quasiconformal mappings". Preprint
- [NTV] F. Nazarov, S. Treil, and A. Volberg "Weak Type Estimates and Cotlar Inequalities for Calderon-Zygmund Operators on Nonhomogeneous Spaces" International Mathematics Research Notices, 1998, #9.
- [P] J. Privalov, "Sur les fonctions conjuguées", Bull. Soc Math. France 44 (1916), 100-103
- [T] X. Tolsa, "BMO, H^1 and Calderon-Zygmund operators for non-doubling measures". Math. Ann. 319 (2001), 89-149.
- [S1] E. M. Stein, Singular Integrals and Differentiability Properties of Functions . Princeton University Press, Princeton N. J.1970
- [S2] E. M. Stein, Harmonic Analysis. Real-Variable Methods, Orthogonality, and Oscillatory Integrals. Princeton University Press. Princeton, New Jersey 1993.
- [W] R. Wittmann, "Application of a Theorem of M. G. Krein to singular Integrals".Transactions of the American Mathematical Society, Vol. 299, #2, February 1987.
- [Z] A. Zygmund, Trigonometric Series, 2nd. ed., Cambridge Univ. Press, Cambridge, 1968

e-mail of the author:
aegatto@depaul.edu